

A NOTE ON THREE TYPES OF QUASISYMMETRIC FUNCTIONS

T. KYLE PETERSEN

ABSTRACT. In the context of generating functions for P -partitions, we revisit three flavors of quasisymmetric functions: Gessel's quasisymmetric functions, Chow's type B quasisymmetric functions, and Poirier's signed quasisymmetric functions. In each case we use the inner coproduct to give a combinatorial description (counting pairs of permutations) to the multiplication in: Solomon's type A descent algebra, Solomon's type B descent algebra, and the Mantaci-Reutenauer algebra, respectively. The presentation is brief and elementary, our main results coming as consequences of P -partition theorems already in the literature.

1. QUASISYMMETRIC FUNCTIONS AND SOLOMON'S DESCENT ALGEBRA

The ring of quasisymmetric functions is well-known (see [8], ch. 7.19). Recall that a quasisymmetric function is a formal series

$$Q(x_1, x_2, \dots) \in \mathbb{Z}[[x_1, x_2, \dots]]$$

of bounded degree such that the coefficient of $x_{i_1}^{\alpha_1} x_{i_2}^{\alpha_2} \cdots x_{i_k}^{\alpha_k}$ is the same for all $i_1 < i_2 < \cdots < i_k$ and all compositions $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_k)$. Recall that a composition of n , written $\alpha \models n$, is an ordered tuple of positive integers $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_k)$ such that $|\alpha| = \alpha_1 + \alpha_2 + \cdots + \alpha_k = n$. In this case we say that α has k parts, or $\#\alpha = k$. We can put a partial order on the set of all compositions of n by reverse refinement. The covering relations are of the form

$$(\alpha_1, \dots, \alpha_i + \alpha_{i+1}, \dots, \alpha_k) \prec (\alpha_1, \dots, \alpha_i, \alpha_{i+1}, \dots, \alpha_k).$$

Let $\mathcal{Qsym}_n$ denote the set of all quasisymmetric functions homogeneous of degree n . The ring of quasisymmetric functions can be defined as $\mathcal{Qsym} := \bigoplus_{n \geq 0} \mathcal{Qsym}_n$, but our focus will be on the quasisymmetric functions of degree n , rather than the ring as a whole.

The most obvious basis for $\mathcal{Qsym}_n$ is the set of *monomial* quasisymmetric functions, defined for any composition $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_k) \models n$,

$$M_\alpha := \sum_{i_1 < i_2 < \dots < i_k} x_{i_1}^{\alpha_1} x_{i_2}^{\alpha_2} \dots x_{i_k}^{\alpha_k}.$$

We can form another natural basis with the *fundamental* quasisymmetric functions, also indexed by compositions,

$$F_\alpha := \sum_{\alpha \preceq \beta} M_\beta,$$

since, by inclusion-exclusion we can express the M_α in terms of the F_α :

$$M_\alpha = \sum_{\alpha \preceq \beta} (-1)^{\#\beta - \#\alpha} F_\beta.$$

As an example,

$$F_{(2,1)} = M_{(2,1)} + M_{(1,1,1)} = \sum_{i < j} x_i^2 x_j + \sum_{i < j < k} x_i x_j x_k = \sum_{i \leq j < k} x_i x_j x_k.$$

Compositions can be used to encode descent classes of permutations in the following way. Recall that a *descent* of a permutation $\pi \in \mathfrak{S}_n$ is a position i such that $\pi_i > \pi_{i+1}$, and that an *increasing run* of a permutation π is a maximal subword of consecutive letters $\pi_{i+1}\pi_{i+2}\dots\pi_{i+r}$ such that $\pi_{i+1} < \pi_{i+2} < \dots < \pi_{i+r}$. By maximality, we have that if $\pi_{i+1}\pi_{i+2}\dots\pi_{i+r}$ is an increasing run, then i is a descent of π (if $i \neq 0$), and $i+r$ is a descent of π (if $i+r \neq n$). For any permutation $\pi \in \mathfrak{S}_n$ define the *descent composition*, $C(\pi)$, to be the ordered tuple listing (from left to right) the lengths of the increasing runs of π . If $C(\pi) = (\alpha_1, \alpha_2, \dots, \alpha_k)$, we can recover the descent set of π :

$$\text{Des}(\pi) = \{\alpha_1, \alpha_1 + \alpha_2, \dots, \alpha_1 + \alpha_2 + \dots + \alpha_{k-1}\}.$$

Since $C(\pi)$ and $\text{Des}(\pi)$ have the same information, we will use them interchangeably. For example, the permutation $\pi = (3, 4, 5, 2, 6, 1)$ has $C(\pi) = (3, 2, 1)$ and $\text{Des}(\pi) = \{3, 5\}$.

Recall ([7], ch. 4.5) that a P -partition is an order-preserving map from a poset P to some (countable) totally ordered set. To be precise, let P be any labeled partially ordered set (with partial order $<_P$) and let S be any totally ordered countable set. Then $f : P \rightarrow S$ is a P -partition if it satisfies the following conditions:

- (1) $f(i) \leq f(j)$ if $i <_P j$
- (2) $f(i) < f(j)$ if $i <_P j$ and $i > j$ (as labels)

We let $\mathcal{A}(P)$ (or $\mathcal{A}(P; S)$ if we want to emphasize the image set) denote the set of all P -partitions, and encode this set in the generating function

$$\Gamma(P) := \sum_{f \in \mathcal{A}(P)} x_{f(1)} x_{f(2)} \cdots x_{f(n)},$$

where n is the number of elements in P (we will only consider finite posets). If we take S to be the set of positive integers, then it should be clear that $\Gamma(P)$ is always going to be a quasisymmetric function of degree n . As an easy example, let P be the poset defined by $3 >_P 2 <_P 1$. In this case we have

$$\Gamma(P) = \sum_{f(3) \geq f(2) < f(1)} x_{f(1)} x_{f(2)} x_{f(3)}.$$

We can consider permutations to be labeled posets with total order $\pi_1 <_\pi \pi_2 <_\pi \cdots <_\pi \pi_n$. With this convention, we have

$$\begin{aligned} \mathcal{A}(\pi) &= \{f : [n] \rightarrow S \mid f(\pi_1) \leq f(\pi_2) \leq \cdots \leq f(\pi_n) \\ &\quad \text{and } k \in \text{Des}(\pi) \Rightarrow f(\pi_k) < f(\pi_{k+1})\}, \end{aligned}$$

and

$$\Gamma(\pi) = \sum_{\substack{i_1 \leq i_2 \leq \cdots \leq i_n \\ k \in \text{Des}(\pi) \Rightarrow i_k < i_{k+1}}} x_{i_1} x_{i_2} \cdots x_{i_n}.$$

It is not hard to verify that in fact we have

$$\Gamma(\pi) = F_{C(\pi)},$$

so that generating functions for the P -partitions of permutations $\pi \in \mathfrak{S}_n$ form a basis for $\mathcal{Qsym}_n$.

We have the following theorem related to P -partitions of permutations, due to Gessel [3].

Theorem 1. *As sets, we have the bijection*

$$\mathcal{A}(\pi; ST) \leftrightarrow \coprod_{\sigma\tau=\pi} \mathcal{A}(\tau; S) \oplus \mathcal{A}(\sigma; T),$$

where ST is the cartesian product of the sets S and T with the lexicographic ordering and $\coprod$ denotes the disjoint union.

Let $X = \{x_1, x_2, \dots\}$ and $Y = \{y_1, y_2, \dots\}$ be two sets of commuting indeterminates. Then we define the bipartite generating function,

$$\Gamma(\pi)(XY) = \sum_{\substack{(i_1, j_1) \leq (i_2, j_2) \leq \cdots \leq (i_n, j_n) \\ k \in \text{Des}(\pi) \Rightarrow (i_k, j_k) < (i_{k+1}, j_{k+1})}} x_{i_1} \cdots x_{i_n} y_{j_1} \cdots y_{j_n}.$$

We will apply Theorem 1 with $S = T = \mathbb{P}$, the positive integers.

Corollary 1. *We have*

$$F_{C(\pi)}(XY) = \sum_{\sigma\tau=\pi} F_{C(\tau)}(X)F_{C(\sigma)}(Y).$$

Following [3], we can define a coalgebra $\mathcal{Qsym}_n^*$ in the following way. If π is any permutation with $C(\pi) = \gamma$, let $a_{\alpha,\beta}^\gamma$ denote the number of pairs of permutations $(\sigma, \tau) \in \mathfrak{S}_n \times \mathfrak{S}_n$ with $C(\sigma) = \alpha$, $C(\tau) = \beta$, and $\sigma\tau = \pi$. Then Corollary 1 defines a coproduct $\mathcal{Qsym}_n^* \rightarrow \mathcal{Qsym}_n^* \otimes \mathcal{Qsym}_n^*$:

$$F_\gamma \mapsto \sum_{\alpha, \beta \models n} a_{\alpha,\beta}^\gamma F_\beta \otimes F_\alpha.$$

The dual space to $\mathcal{Qsym}_n^*$ is then $\mathcal{Qsym}_n$ equipped with multiplication

$$F_\beta * F_\alpha = \sum_{\gamma} a_{\alpha,\beta}^{\gamma \models} F_\gamma.$$

Let $\mathbb{Z}\mathfrak{S}_n$ denote the group algebra of the symmetric group. We can define its dual coalgebra $\mathbb{Z}\mathfrak{S}_n^*$ with comultiplication

$$\pi \mapsto \sum_{\sigma\tau=\pi} \tau \otimes \sigma.$$

Then we have a surjective homomorphism of coalgebras $\varphi^* : \mathbb{Z}\mathfrak{S}_n^* \rightarrow \mathcal{Qsym}_n^*$ given by

$$\varphi^*(\pi) = F_{C(\pi)}.$$

The dualization of this map is then an injective homomorphism $\varphi : \mathcal{Qsym}_n \rightarrow \mathbb{Z}\mathfrak{S}_n$ with

$$\varphi(F_\alpha) = \sum_{C(\pi)=\alpha} \pi.$$

The image of φ is then a subalgebra of the group algebra, with basis

$$u_\alpha := \sum_{C(\pi)=\alpha} \pi.$$

This subalgebra is well-known as *Solomon's descent algebra*, denoted $\text{Sol}(A_{n-1})$. Corollary 1 has then given a combinatorial description to multiplication in $\text{Sol}(A_{n-1})$:

$$u_\beta u_\alpha = \sum_{\gamma \models n} a_{\alpha,\beta}^\gamma u_\gamma.$$

The above arguments are due to Gessel [3]. We give them here in full detail for comparison with later sections, when we will outline a similar relationship between type Chow's B quasisymmetric functions and $\text{Sol}(B_n)$, and between Poirier's *signed* quasisymmetric functions and the Mantaci-Reutenauer algebra.

2. TYPE B QUASISYMMETRIC FUNCTIONS AND SOLOMON'S DESCENT ALGEBRA

The type B quasisymmetric functions can be viewed as the natural objects related to type B P -partitions (see [2]). Define the type B posets (with $2n + 1$ elements) to be posets labeled distinctly by $\{-n, \dots, -1, 0, 1, \dots, n\}$ with the property that if $i <_P j$, then $-j <_P -i$. For example, $-2 >_P 1 <_P 0 <_P -1 >_P 2$ is a type B poset.

Let P be any type B poset, and let $S = \{s_0, s_1, \dots\}$ be any countable totally ordered set with a minimal element s_0 . Then a type B P -partition is any map $f : P \rightarrow \pm S$ such that

- (1) $f(i) \leq f(j)$ if $i <_P j$
- (2) $f(i) < f(j)$ if $i <_P j$ and $i > j$ (as labels)
- (3) $f(-i) = -f(i)$

where $\pm S$ is the totally ordered set

$$\dots < -s_2 < -s_1 < s_0 < s_1 < s_2 < \dots$$

If S is the nonnegative integers, then $\pm S$ is the set of all integers.

The third property of type B P -partitions means that $f(0) = 0$ and the set $\{f(i) \mid i = 1, 2, \dots, n\}$ determines the map f . We let $\mathcal{A}_B(P) = \mathcal{A}_B(P; \pm S)$ denote the set of all type B P -partitions, and define the generating function for type B P -partitions as

$$\Gamma_B(P) := \sum_{f \in \mathcal{A}_B(P)} x_{|f(1)|} x_{|f(2)|} \cdots x_{|f(n)|}.$$

Signed permutations $\pi \in \mathfrak{B}_n$ are type B posets with total order

$$-\pi_n < \dots < -\pi_1 < 0 < \pi_1 < \dots < \pi_n.$$

We then have

$$\begin{aligned} \mathcal{A}_B(\pi) &= \{f : \pm[n] \rightarrow \pm S \mid f(-i) = -f(i), \\ &\quad 0 \leq f(\pi_1) \leq f(\pi_2) \leq \dots \leq f(\pi_n), \\ &\quad \text{and } k \in \text{Des}_B(\pi) \Rightarrow f(\pi_k) < f(\pi_{k+1})\}, \end{aligned}$$

and

$$\Gamma_B(\pi) = \sum_{\substack{0 \leq i_1 \leq i_2 \leq \dots \leq i_n \\ k \in \text{Des}(\pi) \Rightarrow i_k < i_{k+1}}} x_{i_1} x_{i_2} \cdots x_{i_n}.$$

Here, the type B descent set, $\text{Des}_B(\pi)$, keeps track of the ordinary descents as well as a descent in position 0 if $\pi_1 < 0$. Notice that if $\pi_1 < 0$, then $f(\pi_1) > 0$, and $\Gamma_B(\pi)$ has no x_0 terms, as in

$$\Gamma_B((-3, 2, -1)) = \sum_{0 < i \leq j < k} x_i x_j x_k.$$

The possible presence of a descent in position zero is the crucial difference between type A and type B descent sets. Define a *psuedo-composition* of n to be an ordered tuple $\alpha = (\alpha_1, \dots, \alpha_k)$ with $\alpha_1 \geq 0$, and $\alpha_i > 0$ for $i > 1$, such that $\alpha_1 + \dots + \alpha_k = n$. We write $\alpha \Vdash n$ to mean α is a psuedo-composition of n . Define the descent psuedo-composition $C(\pi)$ of a signed permutation π be the lengths of its increasing runs as before, but now we have $\alpha_1 = 0$ if $\pi_1 < 0$. As with ordinary compositions, the partial order on pseudo-compositions of n is given by reverse refinement. We can move back and forth between descent psuedo-compositions and descent sets in exactly the same way as for type A. If $C(\pi) = (\alpha_1, \dots, \alpha_k)$, then we have

$$\text{Des}_B(\pi) = \{\alpha_1, \alpha_1 + \alpha_2, \dots, \alpha_1 + \alpha_2 + \dots + \alpha_{k-1}\}.$$

We will use psuedo-compositions of n to index the type B quasisymmetric functions. Define $\mathcal{BQsym}_n$ as the vector space of functions spanned by the *type B monomial quasisymmetric functions*:

$$M_{B,\alpha} := \sum_{0 < i_2 < \dots < i_k} x_0^{\alpha_1} x_{i_2}^{\alpha_2} \dots x_{i_k}^{\alpha_k},$$

where $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_k)$ is any psuedo-composition of n , or equivalently by the *type B fundamental quasisymmetric functions*:

$$F_{B,\alpha} := \sum_{\alpha \preceq \beta} M_{B,\beta}.$$

The space of all type B quasisymmetric functions is defined as the direct sum $\mathcal{BQsym} := \bigoplus_{n \geq 0} \mathcal{BQsym}_n$. By design, we have

$$\Gamma_B(\pi) = F_{B,C(\pi)}.$$

From Chow [2] we have the following theorem and corollary.

Theorem 2. *As sets, we have the bijection*

$$\mathcal{A}_B(\pi; ST) \leftrightarrow \coprod_{\sigma\tau=\pi} \mathcal{A}_B(\tau; S) \oplus \mathcal{A}_B(\sigma; T).$$

We take $S = T = \mathbb{Z}$ and we have the following.

Corollary 2. *We have*

$$F_{B,C(\pi)}(XY) = \sum_{\sigma\tau=\pi} F_{B,C(\tau)}(X) F_{B,C(\sigma)}(Y).$$

The coalgebra structure on $\mathcal{BQsym}_n$ works just the same as the type A case so we will omit some details. Corollary 2 gives us the coproduct

$$F_{B,\gamma} \mapsto \sum_{\alpha, \beta \Vdash n} b_{\alpha,\beta}^\gamma F_{B,\beta} \otimes F_{B,\alpha},$$

where for any π such that $C(\pi) = \gamma$, $b_{\alpha,\beta}^\gamma$ is the number of pairs of signed permutations (σ, τ) such that $C(\sigma) = \alpha$, $C(\tau) = \beta$, and $\sigma\tau = \pi$. The dual algebra is isomorphic to $\text{Sol}(B_n)$, where if u_α is the sum of all signed permutations with descent psuedo-composition α , the multiplication given by

$$u_\beta u_\alpha = \sum_{\gamma \Vdash n} b_{\alpha,\beta}^\gamma u_\gamma.$$

3. SIGNED QUASISYMMETRIC FUNCTIONS AND THE MANTACI-REUTENAUER ALGEBRA

One thing to have noticed about the generating function for type B P -partitions is that we are losing a certain amount of information when we take absolute values on the subscripts. We can think of signed quasisymmetric functions as arising naturally by dropping this restriction.

For a type B poset P , define the *signed generating function* for type B P -partitions to be

$$\bar{\Gamma}(P) := \sum_{f \in \mathcal{A}_B(P)} x_{f(1)} x_{f(2)} \cdots x_{f(n)},$$

where we will write

$$x_i = \begin{cases} u_i & \text{if } i < 0, \\ v_i & \text{if } i \geq 0. \end{cases}$$

In the case where P is a signed permutation, we have

$$\bar{\Gamma}(\pi) = \sum_{\substack{0 \leq i_1 \leq i_2 \leq \cdots \leq i_n \\ s \in \text{Des}_B(\pi) \Rightarrow i_s < i_{s+1} \\ \pi_s < 0 \Rightarrow x_{i_s} = u_{i_s} \\ \pi_s > 0 \Rightarrow x_{i_s} = v_{i_s}}} x_{i_1} x_{i_2} \cdots x_{i_n},$$

so that now we are keeping track of the set of minus signs of our signed permutation along with the descents. For example,

$$\bar{\Gamma}((-3, 2, -1)) = \sum_{0 < i \leq j < k} u_i v_j u_k.$$

To keep track of both the set of signs and the set of descents, we introduce the *signed compositions* as used in [1]. A signed composition α of n , denoted $\alpha \Vdash n$, is a tuple of nonzero integers $(\alpha_1, \dots, \alpha_k)$ such that $|\alpha_1| + \cdots + |\alpha_k| = n$. For any signed permutation π we will associate a signed composition $sC(\pi)$ by simply recording the length of increasing runs with constant sign, and then recording that sign. For example, if $\pi = (-3, 4, 5, -6, -2, -7, 1)$, then $sC(\pi) = (-1, 2, -2, -1, 1)$. The signed composition keeps track of both the set of signs and the set of descents of the permutation, as we demonstrate with an example.

If $sC(\pi) = (-3, 2, 1, -2, 1)$, then we know that π is a permutation in $\mathfrak{S}_9$ such that π_4, π_5, π_6 , and π_9 are positive, whereas the rest are all negative. The descents of π are in positions 5 and 6. Note that for any ordinary composition of n with k parts, there are 2^k signed compositions, leading us to conclude that there are

$$\sum_{k=1}^n \binom{n-1}{k-1} 2^k = 2 \cdot 3^{n-1}$$

signed compositions of n .

We will use signed compositions to index the signed quasisymmetric functions (see [6]). For any signed composition α , define the *monomial signed quasisymmetric function*

$$\overline{M}_\alpha := \sum_{\substack{i_1 < i_2 < \dots < i_k \\ \alpha_r < 0 \Rightarrow x_{i_r} = u_{i_r} \\ \alpha_r > 0 \Rightarrow x_{i_r} = v_{i_r}}} x_{i_1}^{|\alpha_1|} x_{i_2}^{|\alpha_2|} \dots x_{i_k}^{|\alpha_k|},$$

and the *fundamental signed quasisymmetric function*

$$\overline{F}_\alpha := \sum_{\alpha \preccurlyeq \beta} \overline{M}_\beta.$$

By construction, we have

$$\overline{\Gamma}(\pi) = \overline{F}_{sC(\pi)}.$$

Notice that if we set $u = v$, then our signed quasisymmetric functions become type B quasisymmetric functions.

Let $\mathcal{SQsym}_n$ denote the span of the $\overline{M}_\alpha$ (or $\overline{F}_\alpha$), taken over all $\alpha \Vdash n$. The space of all signed quasisymmetric functions, $\mathcal{SQsym} := \bigoplus_{n \geq 0} \mathcal{SQsym}_n$, is a graded ring whose n -th graded component has rank $2 \cdot 3^{n-1}$.

Theorem 2 is a statement about splitting apart bipartite P -partitions, independent of how we choose to encode the information. So while Corollary 2 is one such way of encoding the information of Theorem 2, the following is another.

Corollary 3. *We have*

$$\overline{F}_{sC(\pi)}(XY) = \sum_{\sigma\tau=\pi} \overline{F}_{sC(\tau)}(X) \overline{F}_{sC(\sigma)}(Y).$$

We define the coalgebra $\mathcal{SQsym}_n^*$ as we did in the earlier cases. Let $\pi \in \mathfrak{B}_n$ be any signed permutation with $sC(\pi) = \gamma$, and let $c_{\alpha,\beta}^\gamma$ be the number of pairs of permutations $(\sigma, \tau) \in \mathfrak{B}_n \times \mathfrak{B}_n$ with $sC(\sigma) = \alpha$,

$sC(\tau) = \beta$, and $\sigma\tau = \pi$. Corollary 3 gives a coproduct $\mathcal{SQsym}_n^* \rightarrow \mathcal{SQsym}_n^* \otimes \mathcal{SQsym}_n^*$:

$$\overline{F}_\gamma \mapsto \sum_{\alpha, \beta \Vdash n} c_{\alpha, \beta}^\gamma \overline{F}_\beta \otimes \overline{F}_\alpha.$$

Multiplication in the dual algebra $\mathcal{SQsym}_n$, the signed quasisymmetric functions of degree n , is given by

$$\overline{F}_\beta * \overline{F}_\alpha = \sum_{\gamma \Vdash n} c_{\alpha, \beta}^\gamma \overline{F}_\gamma.$$

The group algebra of the hyperoctahedral group, $\mathbb{Z}\mathfrak{B}_n$, has a dual coalgebra $\mathbb{Z}\mathfrak{B}_n^*$ with comultiplication given by the map

$$\pi \mapsto \sum_{\sigma\tau=\pi} \tau \otimes \sigma.$$

The following is a surjective homomorphism of coalgebras $\psi^* : \mathbb{Z}\mathfrak{B}_n^* \rightarrow \mathcal{SQsym}_n^*$ given by

$$\psi^*(\pi) = \overline{F}_{sC(\pi)}.$$

The dualization of this map is an injective homomorphism $\psi : \mathcal{SQsym}_n \rightarrow \mathbb{Z}\mathfrak{B}_n$ with

$$\psi(\overline{F}_\alpha) = \sum_{sC(\pi)=\alpha} \pi.$$

The image of ψ is then a subalgebra of $\mathbb{Z}\mathfrak{B}_n$ of dimension $2 \cdot 3^{n-1}$, with basis

$$v_\alpha := \sum_{sC(\pi)=\alpha} \pi.$$

This subalgebra is called the *Mantaci-Reutenauer algebra*, with multiplication given explicitly by

$$v_\beta v_\alpha = \sum_{\gamma \Vdash n} c_{\alpha, \beta}^\gamma v_\gamma.$$

In closing, we remark that this same method is seen in [5], where Stembridge's enriched P -partitions [9] are generalized and put to use to study peak algebras. Variations on the theme can also be found in [4].

REFERENCES

- [1] C. Bonaffé and C. Hohlweg, *Generalized descent algebra and construction of irreducible characters of hyperoctahedral groups*, arXiv: math.CO/0409199 .
- [2] C.-O. Chow, *Noncommutative symmetric functions of type B*, Ph.D. thesis, MIT (2001).
- [3] I. Gessel, *Multipartite P-partitions and inner products of skew Schur functions*, Contemporary Mathematics **34** (1984), 289–317.

- [4] T.K. Petersen, *Cyclic descents and P -partitions*, to appear in Journal of Algebraic Combinatorics.
- [5] T.K. Petersen, *Enriched P -partitions and peak algebras*, arXiv: math.CO/0508041.
- [6] S. Poirier, *Cycle type and descent set in wreath products*, Discrete Mathematics **180** (1998), 315–343.
- [7] R. Stanley, *Enumerative Combinatorics, Volume I*, Wadsworth & Brooks/Cole, 1986.
- [8] R. Stanley, *Enumerative Combinatorics, Volume II*, Cambridge University Press, 2001.
- [9] J. Stembridge, *Enriched P -partitions*, Transactions of the American Mathematical Society **349** (1997), 763–788.